

**Importancia diagnóstica de la
hipofosfatemia persistente en
pacientes pediátricos:
conclusiones de un análisis
retrospectivo y perspectivas del
cribado neonatal**

**Diagnostic significance of
persistent hypophosphatasemia
in pediatric patients – Insights
from a retrospective analysis and
neonatal screening perspectives**

Diagnostic significance of persistent hypophosphatasemia in pediatric patients – Insights from a retrospective analysis and neonatal screening perspectives

María José Muñoz Domene¹, Alberto Suárez Catalina², José María Gómez Vida³, Ricardo Pérez lañez³, Manuel Muñoz Torres^{1,4,5,6}, Cristina García Fontana^{1,4,5,6}, Beatriz García Fontana^{1,4,5,6}

¹Instituto de Investigación Biosanitaria de Granada (Ibs.Granada). Granada. ²Clinical Analysis Unit; ³Pediatric Unit; and ⁴Endocrinology and Nutrition Unit. Hospital Clínico Universitario San Cecilio. Granada. ⁵CIBER of Frailty and Healthy Aging (CIBERFES). Instituto de Salud Carlos III. Madrid. ⁶Department of Medicine. Universidad de Granada. Granada

Received: 02/09/2025

Accepted: 12/11/2025

Correspondence: Cristina García Fontana. Endocrinology and Nutrition Unit. Hospital Clínico Universitario San Cecilio. Av. del Conocimiento, s/n. 18007 Granada
e-mail: cgfontana@fibao.es

Manuel Muñoz Torres y Cristina García Fontana son ambos autores de correspondencia de este artículo.

Funding: This work was supported by Instituto de Salud Carlos III grant (PI21/01069) co-funded by the European Regional Development Fund (FEDER) and by Junta de Andalucía grant (EXC-2023-06). C. G. F., and B. G. F. were funded by grants C1-S0033 and CP22/00022 from the Junta de Andalucía and the Instituto de Salud Carlos III, respectively. The retrospective screening of ALP levels mentioned in this manuscript was funded by Alexion, AstraZeneca Rare Diseases.

Conflict of interest: The authors declare no conflict of interest.

Artificial intelligence: The authors declare not to have used artificial intelligence (AI) or any AI-assisted technologies in the elaboration of the article.

ABSTRACT

Introduction: hypophosphatasia (HPP) is a rare genetic disorder caused by loss-of-function mutations in the *ALPL* gene, which encodes tissue-nonspecific alkaline phosphatase (TNSALP). The resulting enzyme deficiency primarily affects bone and dental mineralization, with clinical manifestations ranging from lethal perinatal forms to milder symptoms in later childhood. Despite the availability of targeted therapies, many cases remain undiagnosed, contributing to significant morbidity and mortality.

Material and methods: we conducted a retrospective study including pediatric patients aged 0-12 years with persistently low alkaline phosphatase (ALP) levels. Medical records were reviewed to exclude secondary causes of hypophosphatasemia. Eligible patients underwent further laboratory testing and radiographic evaluation to assure persistent low ALP levels and to detect hypomineralization consistent with HPP.

Results: out of 271 initially identified patients, 216 were excluded due to secondary causes. Forty-one patients were suspected of having HPP; among these, 23 declined consent for further investigation, 15 could not be contacted and 3 were selected for the screening step of a clinical trial for an investigational HPP treatment. Notably, 12 patients died without a definitive diagnosis of HPP, indicating a concerning mortality rate of 4.4 % in this cohort.

Conclusions: these findings highlight the underdiagnosis of HPP in pediatric practice and underscore the urgent need for early screening strategies, including neonatal detection, to initiate timely treatment and reduce preventable mortality. Education of healthcare providers

and diagnostic algorithms based on serum ALP levels are essential to improve outcomes.

Keywords: Hypophosphatasia. Persistent hypophosphatasemia. ALPL gene. Tissue-nonspecific alkaline phosphatase (TNSALP). Pediatric and neonatal screening.

INTRODUCTION

Hypophosphatasia (HPP) is a rare genetic disorder caused by loss-of-function mutations in the *ALPL* gene, which encodes the tissue-nonspecific isoenzyme of alkaline phosphatase (TNSALP) (1). It was first described by J.C. Rathbun in 1948 (2). This enzyme plays a crucial role in bone and dental mineralization as it hydrolyzes substrates such as inorganic pyrophosphate (PPi), a potent inhibitor of hydroxyapatite crystal formation. Deficiency of TNSALP leads to extracellular accumulation of PPi and other substrates, such as pyridoxal-5'-phosphate (PLP) -circulating form of vitamin B6- and phosphoethanolamine (PEA), impairing normal mineralization (3). The accumulation of PLP should also be considered, as their effects may extend beyond bone metabolism, potentially influencing neurological function. E.g.: impaired hydrolysis of PLP can cause vitamin B6 deficiency in the central nervous system and reduced neurotransmitter synthesis, potentially leading to pyridoxine-responsive seizures in severely affected infants (4).

The first mutation in the *ALPL* gene was identified in 1988 in a Canadian infant with the lethal form of HPP (5). Currently, over 480 mutations linked to HPP have been reported, the majority of which are missense variants (6). HPP can be inherited in either an autosomal recessive or autosomal dominant pattern. The more severe clinical forms, are typically inherited in an autosomal recessive manner. By contrast, milder forms of the disease may result from either autosomal recessive or dominant inheritance (7). The wide

variability in genotype-phenotype correlations raises challenges for clinical prediction and management, especially among heterozygous carriers (8). Importantly, heterozygous individuals can also manifest clinical symptoms, and recent data from the Global HPP Registry indicate that the majority of patients with non-life-threatening forms of HPP carry a single *ALPL* variant (9,10). It has been demonstrated that three categories of alleles, classified according to their severity and dominant effect (s: severe recessive, Sd: severe dominant, and m: moderate), occur with a similar prevalence across different regions of Europe, suggesting that the prevalence of HPP is likely consistent (11). Even though, the current estimate of potential undiagnosed HPP cases in countries like Spain could rise to 27,177 cases (12). This would make the prevalence of HPP higher than suspected.

There is traditionally considered to be an inverse relationship between age at onset and clinical severity in HPP, with the most severe phenotypes typically occurring in the perinatal and infantile forms. However, this concept has evolved as growing evidence shows that adults can also present with substantial disease burden, and in some cases with symptoms that rival or exceed those seen in children. Many adults experience chronic musculoskeletal pain, impaired mobility, recurrent fractures, dental abnormalities, fatigue, and even require assistive devices such as wheelchairs or surgical interventions like intramedullary rodding to stabilize femoral fractures (13,14). The high variability in symptomatology, particularly in adults, contributes to frequent diagnostic delays, often by several years or even decades, as historical classifications have tended to exclude adults with pediatric-onset disease (15). Furthermore, recent data show that HPP is a dynamic and evolving disorder: symptoms can change over time and multiple clinical complications accumulate progressively throughout life, rather than being static or confined to a single body system (13,16). While early-onset forms are associated with high mortality, many patients live into adulthood yet endure a considerable and persistent disease burden, affecting physical

function, daily activities, and overall quality of life (14,17). Consequently, HPP is increasingly understood as a lifelong, multisystemic condition rather than solely a pediatric or bone-related disease, and diagnosis remains based on persistently low serum alkaline phosphatase levels, alongside clinical, radiological, and often genetic findings (18).

Regarding pediatric forms, perinatal HPP is the most severe and typically lethal form of the disease, presenting before or at birth (19,20). It is characterized by profound skeletal hypomineralization, resulting in short, deformed limbs and caput membranaceum. Additional features may include a high-pitched cry, pyridoxine-responsive seizures, episodes of apnea with cyanosis and bradycardia, fever of unknown origin, irritability, bone marrow failure (myelophthistic anemia), intracranial hemorrhage, and occasionally underdeveloped lungs (21,22). Survival is very rare.

Infantile HPP presents before 6 months of age (23). Symptoms such as poor feeding, failure to thrive, and delayed milestones are usually found in these patients (24). It is marked by skeletal abnormalities, metabolic imbalances (like hypercalcemia and hypercalciuria), and complications such as fractures, respiratory issues, and pyridoxine-dependent seizures (25). Thoracic deformities can lead to pneumonia, a major cause of death. Despite initial normal development, the condition often progresses rapidly, with a historically high infant mortality rate (11).

Childhood HPP presents after 6 months of age, and it can be considered mild or severe depending on the clinical presentation (7,26). Premature loss primary teeth is a common feature of childhood HPP, and in severe cases, all deciduous teeth may be lost early (7). Skeletal deformities resembling rickets are often present, including skull abnormalities, bowed legs, knock-knees, and enlarged joints due to metaphyseal flaring. Children may experience significant bone pain, short stature, muscle weakness, delayed walking, and a waddling gait. In rare cases, joint swelling and radiographic changes

may resemble chronic recurrent multifocal osteomyelitis (7,25,27,28). Importantly, this traditional classification based on age of presentation does not fully capture the complexity of the disease, as many adults may have pediatric-onset HPP that persisted undiagnosed for years, while others develop symptoms later in life (adult-onset HPP). Thus, adult patients may belong to either group, underscoring the need to consider disease onset, progression, and cumulative burden rather than age alone for accurate characterization.

Based on this, increased awareness and the implementation of screening protocols are essential for early recognition and management, particularly as enzyme replacement therapy is now available and approved by major regulatory agencies such as the FDA and the EMA (24,29).

METHODS

Study population

A retrospective study was performed in the province of Granada, focusing on the pediatric population with suspected HPP. Patients were identified through the laboratory database by selecting those who had at least one recorded test of low serum ALP levels, defined according to the reference values for the age and sex of the patient (Table I) according to the CALIPER study (30). The electronic medical records of these patients were subsequently review to collect clinical information.

Table I. Reference ranges for serum alkaline phosphatase (ALP) activity by age

Age	ALP reference interval (U/L)
	Female/Male
0 to 14 days	90-273
15 days to < 1 year	134-518

1 to < 10 years	156-369
10 to < 13 years	141-460

Data on age, sex, medical history and usual medication were collected for each patient. All the information was incorporated into an anonymized database for analysis, guaranteeing confidentiality in accordance with current data protection regulations.

From the total number of identified patients, exclusion criteria were applied to avoid secondary causes of HPP since low ALP is not only pathognomonic for this disease but also may be observed in various other conditions (8). Patients were excluded for any of the non-HPP causes of hypophosphatasemia listed in table II.

Table II. Secondary causes of hypophosphatasemia

Clinical Methodological situations	causes - laboratory	Comments/Mechanism
Pernicious or severe anemia		Metabolic disturbances affecting phosphate levels
Celiac disease		Intestinal malabsorption causing phosphate loss
Vitamin C deficiency		Alters bone metabolism and phosphate levels
Zinc or magnesium deficiency		Impairs phosphate absorption and metabolism
Cleidocranial dysplasia		Bone disorder impacting mineral metabolism
Mseleni joint disease		Rare condition with joint and bone involvement affecting phosphate metabolism

Bisphosphonate therapy	Suppresses bone resorption, can decrease serum phosphate
Adynamic renal osteodystrophy	Low bone turnover disease causing phosphate imbalance
Wilson's disease	Liver and kidney damage affecting mineral metabolism
Hypothyroidism	Metabolic changes leading to reduced serum phosphate
Vitamin D intoxication	Increases renal phosphate excretion
Radioactive heavy metals	Renal tubular damage causing phosphate wasting (similar to Fanconi syndrome)
Multiple myeloma	Renal involvement causing tubular phosphate loss
Osteogenesis imperfecta type II	Bone disorder affecting phosphate metabolism
Milk-alkali syndrome	Metabolic alkalosis promoting phosphate shift into cells
Cushing's syndrome	Elevated cortisol increasing renal phosphate excretion
Malnutrition	Reduced intake and absorption of phosphate
Major trauma	Increased cellular uptake and shifts of phosphate
Major surgery	Metabolic stress and redistribution of phosphate
Sepsis	Altered metabolism and redistribution causing hypophosphatemia
Hemochromatosis*	Iron deposition causes renal tubular damage leading to renal phosphate wasting (secondary

	Fanconi syndrome)
Enteropathic acrodermatitis*	Zinc deficiency impairs the activity of zinc-dependent metalloenzymes (including TNSALP)
Blood sample collected improperly (oxalate, EDTA)	Chelating agents artificially lowering measured phosphate levels
Inappropriate reference ranges	May cause misinterpretation of phosphate levels
Clofibrate treatment	Can alter lipid and phosphate metabolism
Massive transfusion	Dilutional and transient biochemical parameter changes
Cardiac bypass surgery	Hemodynamic and metabolic changes affecting phosphate levels

*These causes represent recently identified etiologies of hypophosphatasemia, as reported by Villa-Suárez and Sanabria-De La Torre (31,32).

Potential HPP cases were contacted by their referring specialists or primary care physicians to undergo screening within the framework of the *Phase 3 Study of ALXN1850 in Treatment-Naïve Pediatric Participants with HPP*. As part of the study protocol, these patients were referred to *Hospital Universitario La Paz* for the necessary diagnostic procedures, including genetic sequencing of the *ALPL* gene and determination of serum PLP levels. These patients signed the corresponding informed consent.

The present study was approved by the ethics committee of Granada in accordance with the principles of the World Medical Association Declaration of Helsinki (project ID: 0768-N-17 on 28 June 2017).

RESULTS

A total of 271 pediatric cases aged between 0 and 12 years having at least one recorded blood test indicating low serum ALP levels were identified. The flowchart applied during the screening process is illustrated in figure 1.




Figure 1. Flowchart illustrating patient selection and outcomes. Patients with low serum ALP were first divided into two groups: excluded cases and potential HPP cases. Excluded patients were classified according to the identified secondary causes of hypophosphatasemia. For the potential HPP group, the chart details subsequent follow-up actions and outcomes, including additional diagnostic testing, lack of consent and inability to contact.

Of the 271 patients initially considered, 168 cases (62.0 %) were excluded due to the presence of at least one analytical result showing normal ALP levels during follow-up, suggesting that the initial low value might have been transient or non-specific. Subsequently, 16 patients (5.9 %) were excluded due to a diagnosis of cancer, either under active treatment or within follow-up periods potentially affecting bone metabolism and serum ALP levels. Fifteen patients (5.5 %) had undergone major surgical interventions in the days preceding the detection of low ALP levels, representing another confounding factor, and were therefore excluded.

Additionally, 9 cases (3.3 %) were found to have experienced acute caloric restriction, either due to medical conditions such as severe gastrointestinal disorders or restrictive eating behaviors including anorexia, all of which are known to affect ALP levels and bone turnover. These were also excluded. Five patients (1.8 %) presented with sepsis at the time of laboratory evaluation, which could influence liver and bone enzyme levels. These cases were also discarded from further consideration. Furthermore, isolated cases of pernicious anemia ($n = 1$), celiac disease ($n = 1$), and hypothyroidism ($n = 1$) were identified, each of which constitutes a potential secondary cause of low ALP, and they were therefore removed from the candidate group.

A total of 14 patients (5.2 %) were found to be deceased at the time of data collection, and sufficient follow-up data to evaluate the clinical course or confirmatory diagnostics were unavailable, precluding their inclusion in the final analysis. Their clinical profiles were heterogeneous, and in most cases, complex comorbidities were present (Table III). Firstly, two patients (ages 4 and 3) died of cancer, without evidence suggestive of HPP. Another 4 patients died shortly after being born, associated with perinatal complications such as prematurity, hypoxic-ischemic encephalopathy or congenital

diaphragmatic hernia. In these cases, the underlying cause of HPP remained unexplored due to early death. Eight of the deceased patients exhibited congenital malformations and neurological impairments or their main cause of death was shock, which may be consistent with HPP and therefore could not be definitively excluded as HPP patients.

Table III. Clinical features in death patients

N o.	Age	Sex	Cause of death	Medical conditions
1	5	M	Loss of consciousness	Central nervous system malformation, epilepsy and hip pathology
2	4	M	Cancer	
3	4	F	Respiratory infection	Severe perinatal hypoxic-ischemic encephalopathy, symptomatic epilepsy and microcephaly
4	3	F	Cancer	
5	11 months	F	Died following surgery	Polymalformative syndrome, facial hypoplasia and cleft palate
6	5	M	Cardiopulmonary arrest	Microcephaly and severe psychomotor delay
7	5 months	F	Cardiopulmonary arrest	Timothy syndrome
8	2 days	F	Multiorgan failure	
9	1 month	F	Prematurity	
10	1 month	F	Cardiopulmonary arrest	Severe pulmonary hypertension and congenital diaphragmatic hernia.

11	1 month	M	Prematurity	
12	9 months	F	Cardiopulmonary arrest	Central hypotonia, epileptic encephalopathy, progressive microcephaly, growth retardation, movement disorder with dyskinesias, and non-ketotic hypoglycemia
13	2	F	Exacerbation due to <i>Pseudomonas aeruginosa</i>	Aicardi syndrome, epileptic encephalopathy and chronic respiratory failure
14	1 day	F	Cardiogenic shock due to ventricular dysfunction	Irregular pulmonary aeration with areas of atelectasis and hypoxic-ischemic encephalopathy with mild intraventricular hemorrhage

After applying all exclusion criteria, 41 cases (15.1 %) with no identifiable secondary cause of hypophosphatasemia constituted the group of potential HPP cases, pending further diagnostic workup.

Of the 41 potential HPP cases identified, primary care physicians of 28 patients (68.2 %), were successfully contacted. Among these, three patients underwent additional diagnostic testing. These individuals were evaluated with expanded clinical, biochemical, and radiological assessments, including confirmatory enzymatic studies and, where appropriate, *ALPL* gene sequencing. Another two patients (4.9 %) were identified as foreigners who received isolated medical attention in Andalusian's centers and could not be followed up. Twenty-three families of the patients (56 %) declined consent for further investigations. The remaining 13 primary care physicians

(31.7 %) could not be contacted due to missing or outdated contact information.

Among the three patients (7.3 %) who underwent further diagnostic evaluation, none were ultimately diagnosed with HPP (Table IV). In one case, a repeat blood test showed normal ALP levels, leading to exclusion. The remaining two patients continued to exhibit persistently low ALP levels and underwent bilateral wrist and knee radiographs. One of them showed radiographic signs consistent with rickets, while the other had only mild, nonspecific changes, which were not considered sufficient for HPP diagnosis. The patient with radiographic signs of rickets also presented with short stature and a pectus excavatum deformity; however, subsequent genetic testing for *ALPL* mutations returned negative, thereby excluding the HPP diagnosis.

Table IV. Clinical, biochemical and genetic parameters of selected patients

N o.	Age (years)	Vitamin D levels (reference levels)	Parathyroid hormone (reference levels)	Symptoms	Radiography results	Genetic test
1	8	20.1 ng/mL (13.8-41.0)	-	Pectus excavatum, short stature	Sings of rickets	Negative
2	9	17.2 ng/mL (13.8-41.0)	63.3 pg/mL (36.0-144.0)	Dental problems, frequent respiratory infections	Nonspecific findings	-
3	8	32.3 ng/mL	-	Low weight	Nonspecific	-

		mL (12.0-55.0)			fic findings	
--	--	----------------	--	--	--------------	--

DISCUSSION

This retrospective study aimed to identify potential pediatric cases of HPP based on persistently low serum ALP levels and to assess the clinical consequences of under recognition in this population. Out of 271 children aged 0-12 years with at least one recorded episode of hypophosphatasemia, 41 patients (15.1 %) were identified as having persistently low ALP levels without identifiable secondary causes of hypophosphatasemia. These individuals constitute a group of high clinical interest, underscoring the potential diagnostic value of unexplained hypophosphatasemia in routine pediatric care and highlighting a window of opportunity for earlier identification of HPP.

Among the patients who underwent extended evaluation, two maintained persistently low ALP and one presented with short stature, chest deformity (pectus excavatum), and radiological signs compatible with rickets. Although the clinical and biochemical picture raised strong suspicion of HPP, sequencing of the *ALPL* gene did not identify any pathogenic variants. However, this result must be interpreted with caution, as conventional sequencing methods typically assess only coding regions of the gene and may fail to detect pathogenic variants located in intronic, promoter, or other regulatory non-coding regions (33,34). In such cases, functional *ALPL* gene expression may be impaired despite the absence of detectable coding mutations. Therefore, a diagnosis of HPP cannot be definitively excluded in this patient, reflecting the diagnostic limitations in clinical practice, particularly in patients with borderline or atypical phenotypes. According to the article by Rush et al., patients may be diagnosed with HPP not only when pathogenic or likely pathogenic variants in the *ALPL* gene are identified, but also when variants of uncertain significance (VUS) are present, or even in the absence of

detectable variants, provided that the clinical and biochemical features are consistent with the disease (35). In these cases, additional assessments are recommended to be performed such as whole exome sequencing to identify potential mutations in non-coding regions of *ALPL* gene or in other ALP-regulatory genes.

Despite the rigorous inclusion criteria and systematic screening approach, no confirmed cases of HPP were diagnosed in this cohort. This finding is consistent considering the reported prevalence of HPP in its more severe forms. Estimates suggest that the combined prevalence of perinatal and infantile HPP (typically inherited in an autosomal recessive pattern), ranges from 1 in 100,000 to 1 in 300,000 live births (11,12). Therefore, given the limited sample size and geographic focus of this study, the absence of confirmed diagnoses aligns with statistical expectations for severe forms. Additionally, a significant proportion of potentially eligible patients (31.7 %) could not be evaluated due to loss of contact with primary care physicians or refusal of consent by the families. This represents a substantial diagnostic gap that may have masked the identification of true HPP cases, especially among patients with milder or atypical presentations. These logistical limitations, which are common in retrospective studies involving rare diseases, highlight the challenges of case detection in real-world clinical settings.

One of the most concerning findings of this study was the identification of 14 pediatric patients who died before further evaluation could be performed. Only two of these (patients 2 and 4) had a clearly documented oncologic cause of death, with no clinical suspicion of HPP. In the remaining 12 patients, HPP could not be ruled out as a contributing or primary cause of death. Notably, several of these individuals presented with clinical features that may be consistent with severe forms of HPP, including early-onset neurological impairment, epileptic encephalopathy, congenital malformations, growth delay, and severe respiratory dysfunction. Patients 1, 3, 12, and 13 had epilepsy or epileptic encephalopathy,

manifestations that are potentially linked to PLP accumulation in the context of TNSALP deficiency, as the enzyme plays a key role in the dephosphorylation of PLP to pyridoxal, allowing its cellular and central nervous system uptake (36). PLP accumulation may disrupt neurotransmitter synthesis, contributing to seizure activity (37).

Additionally, patients 3, 6, 7, 10, 12, 13, and 14 died from cardiorespiratory arrest or respiratory complications. Patient 14 had documented pulmonary atelectasis, a finding that may be mechanistically linked to impaired phospholipid biosynthesis in the setting of TNSALP deficiency, although rib cage deformities and hypoplastic lungs are more commonly recognized as contributing factors in HPP-related respiratory failure (38,39). Furthermore, multiple patients presented with syndromic phenotypes involving facial or skeletal malformations (e.g., patients 5, 12, and 13), which could align with the perinatal or infantile forms of HPP that are often lethal if not recognized and treated early (21).

The mortality rate observed in this cohort (4.4 %), excluding oncologic causes, should be interpreted with caution, as the retrospective design and substantial loss to follow-up limit the strength of the conclusions. Nonetheless, this finding underscores a potentially important diagnostic gap in pediatric HPP. Several unexplored deaths occurred in children with persistent hypophosphatasemia, whose reported symptoms closely resembled the classical presentation of HPP. While a direct causal relationship cannot be established, these observations raise concern that some fatal cases might have been related to undiagnosed HPP. This highlights the critical importance of early recognition and diagnostic evaluation in pediatric patients with persistently low alkaline phosphatase levels, as timely identification could prevent severe disease progression and potentially avoid fatal outcomes.

These findings underscore the potential value of developing structured diagnostic algorithm that incorporates ALP as a frontline screening biomarker, particularly when levels are markedly low and

not attributable to secondary conditions. In any child with persistently low ALP and signs of poor mineralization, delayed milestones, or unexplained seizures, HPP should be included in the differential diagnosis. Repeat testing to confirm ALP persistence and exclusion of confounding factors are essential steps, followed by targeted biochemical and genetic testing.

Moreover, our results support the implementation of neonatal biochemical screening strategies for early detection of HPP. Although widespread genetic screening via the heel-prick test remains financially and logistically unfeasible, a more accessible and cost-effective approach could involve measuring ALP activity in capillary blood samples collected shortly after birth. Such a biochemical screening tool, if incorporated into routine newborn testing, could help identify neonates with severe TNSALP deficiency before the onset of life-threatening symptoms. Even though, further studies are required to assess its feasibility, cost-effectiveness, and clinical utility. Infants flagged by this method could then undergo confirmatory testing and early intervention, including consideration for enzyme replacement therapy with asfotase alfa, which has demonstrated significant survival benefits in perinatal and infantile HPP (20,40).

Despite its limitations, including its retrospective nature, reliance on existing laboratory records, and the inability to follow up with a significant proportion of potential cases, this study provides valuable insights into the magnitude of underdiagnosis in pediatric HPP. A major limitation was the difficulty in contacting primary care providers, compounded by time constraints and the high workload within primary care settings, which likely contributed to missed follow-up opportunities. Furthermore, the lack of access to advanced genetic or enzymatic diagnostic tools across the healthcare system limits clinicians' ability to fully characterize or confirm suspected cases (41). Nevertheless, this study has notable strengths. It is, to our knowledge, one of the first to apply a systematic screening approach based on real-world ALP data to a pediatric population, offering a

practical framework for identifying children who may benefit from further diagnostic workup. The clinical data gathered from deceased patients highlights the potentially severe consequences of undiagnosed HPP and demonstrates how routine biochemical markers, if correctly interpreted, can uncover life-threatening conditions that might otherwise remain hidden. By focusing on the pediatric population, this study addresses the group at highest risk for the severe, life-limiting forms of the disease and helps to define the profile of patients who would benefit most from early detection and intervention.

In conclusion, this study highlights the importance of persistent hypophosphatasemia as an early warning sign of HPP in children and underscores the substantial mortality burden that may go unrecognized in the absence of structured diagnostic protocols. The development of standardized clinical algorithms and the implementation of biochemical neonatal screening strategies are critical next steps to improve diagnostic yield and reduce avoidable morbidity and mortality in pediatric HPP. In the era of effective enzyme replacement therapy, early identification is no longer optional, it is essential.

REFERENCES

1. Mornet E. Hypophosphatasia. Orphanet J Rare Dis. 4 de octubre de 2007;2:40. DOI: 10.1186/1750-1172-2-40
2. Rathbun JC. Hypophosphatasia; a new developmental anomaly. Am J Dis Child (1911) 1948;75(6):822-31. DOI: 10.1001/archpedi.1948.02030020840003
3. Whyte MP, Mahuren JD, Vrabel LA, Coburn SP. Markedly increased circulating pyridoxal-5'-phosphate levels in hypophosphatasia. Alkaline phosphatase acts in vitamin B6 metabolism. J Clin Invest 1985;76(2):752-6. DOI: 10.1172/JCI112031

4. Baumgartner-Sigl S, Haberlandt E, Mumm S, Scholl-Bürgi S, Sergi C, Ryan L, et al. Pyridoxine-responsive seizures as the first symptom of infantile hypophosphatasia caused by two novel missense mutations (c.677T>C, p.M226T; c.1112C>T, p.T371I) of the tissue-nonspecific alkaline phosphatase gene. *Bone* 2007;40(6):1655-61. DOI: 10.1016/j.bone.2007.01.020
5. Henthorn PS, Whyte MP. Missense mutations of the tissue-nonspecific alkaline phosphatase gene in hypophosphatasia. *Clin Chem* 1992;38(12):2501-5.
6. Farman MR, Rehder C, Malli T, Rockman-Greenberg C, Dahir K, Martos-Moreno GÁ, et al. The Global ALPL gene variant classification project: Dedicated to deciphering variants. *Bone* 2024;178:116947. DOI: 10.1016/j.bone.2023.116947
7. Whyte MP, Zhang F, Wenkert D, McAlister WH, Mack KE, Benigno MC, et al. Hypophosphatasia: Validation and expansion of the clinical nosology for children from 25years experience with 173 pediatric patients. *Bone* 2015;75:229-39. DOI: 10.1016/j.bone.2015.02.022
8. Im M, Cho SY. Hypophosphatasia in childhood: Diagnosis to management. *Osteoporos Sarcopenia* 2025;11(2):38-42. DOI: 10.1016/j.afos.2025.05.003
9. Kishnani PS, Seefried L, Dahir KM, Martos-Moreno GÁ, Högler W, Greenberg CR, et al. Disease burden by *ALPL* variant number in patients with non-life-threatening hypophosphatasia in the Global HPP Registry. *J Med Genet* 2025;62(4):249-57. DOI: 10.1136/jmg-2024-110383
10. Kishnani PS, Seefried L, Dahir KM, Martos-Moreno GÁ, Linglart A, Petryk A, et al. New insights into the landscape of *ALPL* gene variants in patients with hypophosphatasia from the Global HPP Registry. *American J of Med Genetics Pt A* 2024;194(11):e63781. DOI: 10.1002/ajmg.a.63781
11. Mornet E, Taillandier A, Domingues C, Dufour A, Benaloun E, Lavaud N, et al. Hypophosphatasia: a genetic-based nosology and

- new insights in genotype-phenotype correlation. *Eur J Hum Genet* 2021;29(2):289-99. DOI: 10.1038/s41431-020-00732-6
12. González-Cejudo T, Villa-Suárez JM, Ferrer-Millán M, Andújar-Vera F, Contreras-Bolívar V, Andreo-López MC, et al. Mild hypophosphatasia may be twice as prevalent as previously estimated: an effective clinical algorithm to detect undiagnosed cases. *Clinical Chemistry and Laboratory Medicine (CCLM)* 2024;62(1):128-37. DOI: 10.1515/cclm-2023-0427
13. Szabo SM, Tomazos IC, Petryk A, Powell LC, Donato BMK, Zarate YA, et al. Frequency and age at occurrence of clinical manifestations of disease in patients with hypophosphatasia: a systematic literature review. *Orphanet J Rare Dis* 2019;14(1):85. DOI: 10.1186/s13023-019-1062-0
14. Seefried L, Dahir K, Petryk A, Höglér W, Linglart A, Martos-Moreno GÁ, et al. Burden of Illness in Adults With Hypophosphatasia: Data From the Global Hypophosphatasia Patient Registry. *J Bone Miner Res* 2020;35(11):2171-8. DOI: 10.1002/jbmr.4130
15. Höglér W, Langman C, Gomes Da Silva H, Fang S, Linglart A, Ozono K, et al. Diagnostic delay is common among patients with hypophosphatasia: initial findings from a longitudinal, prospective, global registry. *BMC Musculoskelet Disord* 2019;20(1):80. DOI: 10.1186/s12891-019-2420-8
16. Dahir KM, Seefried L, Kishnani PS, Petryk A, Höglér W, Linglart A, et al. Clinical profiles of treated and untreated adults with hypophosphatasia in the Global HPP Registry. *Orphanet J Rare Dis* 2022;17(1):277. DOI: 10.1186/s13023-022-02393-8
17. Dahir KM, Kishnani PS, Martos-Moreno GÁ, Linglart A, Petryk A, Rockman-Greenberg C, et al. Impact of muscular symptoms and/or pain on disease characteristics, disability, and quality of life in adult patients with hypophosphatasia: A cross-sectional analysis from the Global HPP Registry. *Front Endocrinol* 2023;14:1138599. DOI: 10.3389/fendo.2023.1138599

18. Araci MB, Akgun B, Atik T, Isik E, Ak G, Barutcuoglu B, et al. Clinical and molecular findings in children and young adults with persistent low alkaline phosphatase concentrations. *Ann Clin Biochem* 2021;58(4):335-41. DOI: 10.1177/00045632211000102
19. Leung ECW, Mhanni AA, Reed M, Whyte MP, Landy H, Greenberg CR. Outcome of perinatal hypophosphatasia in manitoba mennonites: a retrospective cohort analysis. *JIMD Rep* 2013;11:73-8. DOI: 10.1007/8904_2013_224
20. Whyte MP, Rockman-Greenberg C, Ozono K, Riese R, Moseley S, Melian A, et al. Asfotase Alfa Treatment Improves Survival for Perinatal and Infantile Hypophosphatasia. *J Clin Endocrinol Metab* 2016;101(1):334-42. DOI: 10.1210/jc.2015-3462
21. Shohat M, Rimoin DL, Gruber HE, Lachman RS. Perinatal lethal hypophosphatasia; clinical, radiologic and morphologic findings. *Pediatr Radiol* 1991;21(6):421-7. DOI: 10.1007/BF02026677
22. Silver MM, Vilos GA, Milne KJ. Pulmonary hypoplasia in neonatal hypophosphatasia. *Pediatr Pathol* 1988;8(5):483-93. DOI: 10.3109/15513818809022304
23. Whyte MP. Hypophosphatasia. En: *Genetics of Bone Biology and Skeletal Disease* [Internet]. Elsevier; 2013 [citado 1 de julio de 2025]. p. 337-60. DOI: 10.1016/B978-0-12-387829-8.00022-6. Disponible en: <https://linkinghub.elsevier.com/retrieve/pii/B9780123878298000226>
24. Whyte MP, Greenberg CR, Salman NJ, Bober MB, McAlister WH, Wenkert D, et al. Enzyme-Replacement Therapy in Life-Threatening Hypophosphatasia. *N Engl J Med* 2012;366(10):904-13. DOI: 10.1056/NEJMoa1106173
25. Collmann H, Mornet E, Gattenlöhner S, Beck C, Girschick H. Neurosurgical aspects of childhood hypophosphatasia. *Childs Nerv Syst* 2009;25(2):217-23. DOI: 10.1007/s00381-008-0708-3
26. Fallon MD, Teitelbaum SL, Weinstein RS, Goldfischer S, Brown DM, Whyte MP. Hypophosphatasia: clinicopathologic comparison of the

- infantile, childhood, and adult forms. *Medicine (Baltimore)* 1984;63(1):12-24.
27. Seshia SS, Derbyshire G, Haworth JC, Hoogstraten J. Myopathy with hypophosphatasia. *Arch Dis Child* 1990;65(1):130-1. DOI: 10.1136/adc.65.1.130
28. Whyte MP, Wenkert D, McAlister WH, Mughal MZ, Freemont AJ, Whitehouse R, et al. Chronic recurrent multifocal osteomyelitis mimicked in childhood hypophosphatasia. *J Bone Miner Res* 2009;24(8):1493-505. DOI: 10.1359/jbmr.090308
29. Kishnani PS, Rockman-Greenberg C, Rauch F, Bhatti MT, Moseley S, Denker AE, et al. Five-year efficacy and safety of asfotase alfa therapy for adults and adolescents with hypophosphatasia. *Bone* 2019;121:149-62. DOI: 10.1016/j.bone.2018.12.011
30. Colantonio DA, Kyriakopoulou L, Chan MK, Daly CH, Brinc D, Venner AA, et al. Closing the Gaps in Pediatric Laboratory Reference Intervals: A CALIPER Database of 40 Biochemical Markers in a Healthy and Multiethnic Population of Children. *Clinical Chemistry* 2012;58(5):854-68. DOI: 10.1373/clinchem.2011.177741
31. Villa-Suárez JM, García-Fontana C, Andújar-Vera F, González-Salvatierra S, De Haro-Muñoz T, Contreras-Bolívar V, et al. Hypophosphatasia: A Unique Disorder of Bone Mineralization. *IJMS* 2021;22(9):4303. DOI: 10.3390/ijms22094303
32. Sanabria-de La Torre R, Martínez-Heredia L, González-Salvatierra S, Andújar-Vera F, Iglesias-Baena I, Villa-Suárez JM, et al. Characterization of Genetic Variants of Uncertain Significance for the ALPL Gene in Patients with Adult Hypophosphatasia. *Front Endocrinol* 2022;13:863940. DOI: 10.3389/fendo.2022.863940
33. García-Fontana C, Villa-Suárez JM, Andújar-Vera F, González-Salvatierra S, Martínez-Navajas G, Real PJ, et al. Epidemiological, Clinical and Genetic Study of Hypophosphatasia in A Spanish Population: Identification of Two Novel Mutations in The Alpl Gene. *Sci Rep* 2019;9(1):9569. DOI: 10.1038/s41598-019-46004-2

34. Martínez-Heredia L, Muñoz-Torres M, Sanabria-de La Torre R, Jiménez-Ortas Á, Andújar-Vera F, González-Cejudo T, et al. Systemic effects of hypophosphatasia characterization of two novel variants in the ALPL gene. *Front Endocrinol* 2024;14:1320516. DOI: 10.3389/fendo.2023.1320516
35. Rush E, Brandi ML, Khan A, Ali DS, Al-Alwani H, Almonaei K, et al. Proposed diagnostic criteria for the diagnosis of hypophosphatasia in children and adolescents: results from the HPP International Working Group. *Osteoporos Int* 2024;35(1):1-10. DOI: 10.1007/s00198-023-06843-2
36. Salles JP. Hypophosphatasia: Biological and Clinical Aspects, Avenues for Therapy. *CBR* 2020;41(1):13-27. DOI: 10.33176/AACB-19-00031
37. Waymire KG, Mahuren JD, Jaje JM, Guilarte TR, Coburn SP, MacGregor GR. Mice lacking tissue non-specific alkaline phosphatase die from seizures due to defective metabolism of vitamin B-6. *Nat Genet* 1995;11(1):45-51. DOI: 10.1038/ng0995-45
38. Liedtke D, Hofmann C, Jakob F, Klopocki E, Graser S. Tissue-Nonspecific Alkaline Phosphatase-A Gatekeeper of Physiological Conditions in Health and a Modulator of Biological Environments in Disease. *Biomolecules* 2020;10(12):1648. DOI: 10.3390/biom10121648
39. Padidela R, Yates R, Benscoter D, McPhail G, Chan E, Nichani J, et al. Characterization of tracheobronchomalacia in infants with hypophosphatasia. *Orphanet J Rare Dis* 2020;15(1):204. DOI: 10.1186/s13023-020-01483-9
40. Whyte MP, Madson KL, Phillips D, Reeves AL, McAlister WH, Yakimoski A, et al. Asfotase alfa therapy for children with hypophosphatasia. *JCI Insight* 2016;1(9):e85971. DOI: 10.1172/jci.insight.85971
41. Whyte MP, McAlister WH, Mack KE, Mumm S, Madson KL. Pediatric hypophosphatasia: avoid diagnosis missteps! *J Bone and Miner Res* 2024;39(6):655-60. DOI: 10.1093/jbmr/zjae098

